arXiv:math-ph/0607013v4 2 Dec 2006

A variational formulation of analytical mechanics in an affine space

Antonio De Nicola
Dipartimento di Matematica, Università di Bari
Via Orabona, 4 – 70125 Bari, Italy
`adenicola@tin.it`

Włodzimierz M. Tulczyjew
Associated with
Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Italy
Valle San Benedetto, 2 – 62030 Montecavallo (MC), Italy
`tulczy@libero.it`

Abstract

Variational formulations of statics and dynamics of mechanical systems controlled by external forces are presented as examples of variational principles.

2000 Mathematics Subject Classification: 70G75, 49S05, 70C20, 70H03, 70H05.
Keywords: variational principles, constitutive set, Lagrangian mechanics, Hamiltonian mechanics, Legendre transformation.

1. Introduction.

Two examples of variational principles following the general framework proposed in [1] are presented. The first example is a formulation of statics of mechanical systems. In this formulation the response of a system to control by external forces and not just equilibrium configurations is studied. Information on this response is contained in the constitutive set. The concept of a constitutive set is well known in thermostatics but not usually used in mechanics. The example of statics is presented as the simplest model of a variational principle. The second example is that of dynamics of an autonomous mechanical system. The objects needed in the formulation of a variational principle are tangent vectors and cotangent vectors in a configuration space (as in statics) or a space of motions (in dynamics). In the case of statics the configuration space is a manifold and vectors and covectors are well defined. The space of motions is not a manifold. We construct in this space enough differential structure to permit the construction of vectors and covectors. The construction is a simple example of similar constructions needed for variational formulations of field theory.

A partial presentation of a variational formulation of mechanics was given in [2] within the framework of the geometry of differential manifolds. Affine geometry is used in the present note. This permits a more complete and comprehensible presentation of variational principles.

2. Virtual work principle in statics.

Let Q be an affine space modelled on a vector space V. We consider a static mechanical system with Q as its *configuration space.* Let

$$U: Q \to \mathbb{R} \tag{1}$$

be the *internal energy of the system.* The derivative of U is the mapping

$$\mathrm{D}U: Q \times V \to \mathbb{R}: (q, \delta q) \mapsto \frac{\mathrm{d}}{\mathrm{d}s} U(q + s\delta q)\big|_{s = 0} \tag{2}$$

linear in its vector argument δq. The behaviour of the system controlled by external forces $f \in V^*$ is described by the *constitutive set*

$$S = \Big\{(q, f) \in Q \times V^*;\ \forall_{\delta q \in V}\ \langle f, \delta q\rangle = \mathrm{D}U(q, \delta q)\Big\}. \tag{3}$$

This definition of the constitutive set is the simplest case of a *variational principle* known as the *principle of virtual work.* We will use it as a model for variational principles in analytical mechanics and, in subsequent papers, in electrodynamics and general field theories.

We will now present an example of a well known system to illustrate the concept of the constitutive set.

Example 1. The configuration space Q of a material point of mass m is the affine physical space of three dimensions modelled on a vector space V endowed with an Euclidean metric $g: V \to V^*$. Let the material point $q \in Q$ be connected with a spring of spring constant k to a fixed point $q_0 \in Q$.

The function

$$U: Q \to \mathbb{R}: q \mapsto \frac{k}{2}\langle g(q - q_0), q - q_0\rangle \tag{4}$$

represents the internal energy of the material point. The constitutive set of the system controlled by external forces $f \in V^*$ is the set

$$\begin{aligned} S &= \Big\{(q, f) \in Q \times V^*;\ \forall_{\delta q \in V}\ k\langle g(q - q_0), \delta q\rangle - \langle f, \delta q\rangle = 0\Big\} \\ &= \big\{(q, f) \in Q \times V^*;\ f = kg(q - q_0)\big\}. \end{aligned} \tag{5}$$

The first line of the formula is a variational definition of the constitutive set. ▲

The example describes the statics of the harmonic oscillator and the mass is irrelevant. The dynamics of the same system will be described in a later example.

3. Virtual action principle in analytical mechanics in a finite time interval.

Let the configuration space of an autonomous system be the affine space Q. A *motion* of the system is a parameterized arc in the configuration space – a mapping of an interval $[t_0, t_1]$ in Q differentiable in the sense that it is the restriction of a differentiable mapping defined in an open interval $I \supset [t_0, t_1]$ to the interval $[t_0, t_1]$. The space of motions associated with an interval $[t_0, t_1]$ will be denoted by $\Xi(Q|\mathbb{R}, [t_0, t_1])$ and the space of all motions will be denoted by $\Xi(Q|\mathbb{R})$. A motion $\xi\colon [t_0, t_1] \to Q$ will be denoted by $(\xi|[t_0, t_1])$.

From a differentiable function

$$\kappa\colon Q \times V \times \mathbb{R} \to \mathbb{R} \tag{6}$$

with arguments $q \in Q$, $\dot{q} \in V$, and $t \in \mathbb{R}$ we construct the function

$$K\colon \Xi(Q|\mathbb{R}) \to \mathbb{R}\colon (\xi|[t_0, t_1]) \mapsto \int_{t_0}^{t_1} \kappa \circ (\xi, \dot{\xi}, t). \tag{7}$$

Here the mapping $\dot{\xi}$ is the *velocity* along the motion $\xi\colon [t_0, t_1] \to Q$ defined by

$$\dot{\xi}\colon [t_0, t_1] \to V\colon t \mapsto \frac{\mathrm{d}\xi(t)}{\mathrm{d}t} \tag{8}$$

and $(\xi, \dot{\xi}, t)$ is the mapping

$$(\xi, \dot{\xi}, t)\colon [t_0, t_1] \to Q \times V \times \mathbb{R} : t \mapsto (\xi(t), \dot{\xi}(t), t). \tag{9}$$

The symbol $\Phi(Q|\mathbb{R})$ will be used to denote the space of functions on $\Xi(Q|\mathbb{R})$ constructed in this way from functions on $Q \times V \times \mathbb{R}$.

The space $\Xi(Q|\mathbb{R}, [t_0, t_1])$ is an infinite dimensional affine space. Its model is the vector space of mappings

$$\delta\xi\colon [t_0, t_1] \to V \tag{10}$$

defined on the interval $[t_0, t_1]$. This space of *displacements* will be denoted by $\Delta(Q|\mathbb{R}, [t_0, t_1])$. The space of all displacements is the union

$$\Delta(Q|\mathbb{R}) = \bigcup_{[t_0, t_1] \subset \mathbb{R}} \Delta(Q|\mathbb{R}, [t_0, t_1]). \tag{11}$$

There is no obvious choice of the space dual to $\Delta(Q|\mathbb{R}, [t_0, t_1])$. We use the derivatives of functions in $\Phi(Q|\mathbb{R})$ as models of covectors. The derivative $\mathrm{D}K$ of a function K is defined for each interval $[t_0, t_1]$ separately. It is evaluated on a motion $(\xi|[t_0, t_1]) \in \Xi(Q|\mathbb{R}, [t_0, t_1])$

and a displacement $(\delta\xi|[t_0,t_1]) \in \Delta(Q|\mathbb{R},[t_0,t_1])$. The result is the expression

$$\begin{aligned}
\mathrm{D}K((\xi|[t_0,t_1]),(\delta\xi|[t_0,t_1])) &= \frac{\mathrm{d}}{\mathrm{d}s}K(\xi+s\delta\xi|[t_0,t_1])\big|_{s=0} \\
&= \int_{t_0}^{t_1} \frac{\mathrm{d}}{\mathrm{d}s}\kappa\circ(\xi+s\delta\xi,\dot\xi+s\delta\dot\xi,t)\big|_{s=0} \\
&= \int_{t_0}^{t_1} \left(\left\langle\frac{\partial\kappa}{\partial q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle + \left\langle\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t),\delta\dot\xi\right\rangle\right) \\
&= \int_{t_0}^{t_1} \left(\left\langle\frac{\partial\kappa}{\partial q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle - \left\langle\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t)\right),\delta\xi\right\rangle\right. \\
&\qquad \left.+\frac{\mathrm{d}}{\mathrm{d}t}\left\langle\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle\right) \\
&= \int_{t_0}^{t_1} \left(\left\langle\frac{\partial\kappa}{\partial q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle - \left\langle\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t)\right),\delta\xi\right\rangle\right) \\
&\qquad + \left\langle\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle(t_1) - \left\langle\frac{\partial\kappa}{\partial\dot q}\circ(\xi,\dot\xi,t),\delta\xi\right\rangle(t_0).
\end{aligned} \tag{12}$$

We are using partial derivatives

$$\frac{\partial\kappa}{\partial q}\colon Q\times V\times\mathbb{R}\to V^* \tag{13}$$

defined by

$$\left\langle\frac{\partial\kappa}{\partial q}(q,\dot q,t),\delta q\right\rangle = \frac{\mathrm{d}}{\mathrm{d}s}\kappa(q+s\delta q,\dot q,t)\big|_{s=0} \tag{14}$$

and

$$\frac{\partial\kappa}{\partial\dot q}\colon Q\times V\times\mathbb{R}\to V^* \tag{15}$$

defined by

$$\left\langle\frac{\partial\kappa}{\partial\dot q}(q,\dot q,t),\delta\dot q\right\rangle = \frac{\mathrm{d}}{\mathrm{d}s}\kappa(q,\dot q+s\delta\dot q,t)\big|_{s=0}. \tag{16}$$

The symbol $\delta\dot\xi$ denotes the mapping defined by

$$\delta\dot\xi\colon[t_0,t_1]\to V\colon t\mapsto\frac{\mathrm{d}\delta\xi(t)}{\mathrm{d}t} \tag{17}$$

and $\delta\dot q$ is a generic element of V. The expression (12) for the derivative suggests that a covector is represented by a triple

$$(\varphi|[t_0,t_1],p_0,p_1)\in\Sigma(V^*|\mathbb{R},[t_0,t_1])\times V^*\times V^*, \tag{18}$$

where the symbol $\Sigma(V^*|\mathbb{R}, [t_0, t_1])$ denotes the space of the differentiable mappings of $[t_0, t_1]$ in V^*. The space

$$\Pi(Q; \mathbb{R}|[t_0, t_1]) = \Sigma(V^*|\mathbb{R}, [t_0, t_1]) \times V^* \times V^* \tag{19}$$

with the pairing

$$\langle\!\langle(\varphi|[t_0, t_1], p_0, p_1), (\delta\xi|[t_0, t_1])\rangle\!\rangle = -\int_{t_0}^{t_1} \langle \varphi, \delta\xi \rangle + \langle p_1, \delta\xi(t_1) \rangle - \langle p_0, \delta\xi(t_0) \rangle \tag{20}$$

will be used as the dual of $\Delta(Q|\mathbb{R}, [t_0, t_1])$. The space of all covectors is the union

$$\Pi(Q|\mathbb{R}) = \bigcup_{[t_0, t_1] \subset \mathbb{R}} \Pi(Q|\mathbb{R}, [t_0, t_1]). \tag{21}$$

This construction of the dual spaces is a consequence of the choice of the space $\Phi(Q|\mathbb{R})$ of functions on $\Xi(Q|\mathbb{R})$ and the choice is dictated by the fact that the *action* for a mechanical system is typically one of such functions. The component φ is the *external force* applied to the system. The components p_0 and p_1 are the *initial momentum* and the *final momentum* respectively.

We will formulate the *principle of virtual action* for an autonomous mechanical system. Let

$$L \colon Q \times V \to \mathbb{R} \tag{22}$$

be the *Lagrangian* of the system. The *action* is the function

$$W \colon \Xi(Q|\mathbb{R}) \to \mathbb{R} \colon (\xi|[t_0, t_1]) \mapsto \int_{t_0}^{t_1} L \circ (\xi, \dot{\xi}). \tag{23}$$

The principle of virtual action is included in the definition of the *dynamics*. The *dynamics in the interval* $[t_0, t_1]$ is the set

$$\begin{aligned} D_{[t_0,t_1]} = \Big\{ & ((\xi|[t_0, t_1]), (\varphi|[t_0, t_1], p_0, p_1)) \in \Xi(Q|\mathbb{R}, [t_0, t_1]) \times \Pi(Q; \mathbb{R}|[t_0, t_1]); \\ & \forall_{(\delta\xi|[t_0,t_1]) \in \Delta(Q|\mathbb{R},[t_0,t_1])} \; \langle\!\langle(\varphi|[t_0, t_1], p_0, p_1), (\delta\xi|[t_0, t_1])\rangle\!\rangle = \mathrm{D}W(\xi, \delta\xi|[t_0, t_1]) \Big\}. \end{aligned} \tag{24}$$

The set $D_{[t_0,t_1]}$ corresponds to the constitutive set S used in statics. The space of motions $\Xi(Q|\mathbb{R})$ has in dynamics a role similar to that played in statics by the configuration space Q.

The *dynamics* is the family

$$D = \bigcup_{[t_0, t_1] \subset \mathbb{R}} D_{[t_0,t_1]}. \tag{25}$$

The equality

$$\begin{aligned}\int_{t_0}^{t_1} \left\langle \frac{\partial L}{\partial q} \circ (\xi, \dot{\xi}) - \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}) \right), \delta\xi \right\rangle + \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}), \delta\xi \right\rangle (t_1) \\ - \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}), \delta\xi \right\rangle (t_0) = - \int_{t_0}^{t_1} \langle \varphi, \delta\xi \rangle + \langle p_1, \delta\xi(t_1) \rangle - \langle p_0, \delta\xi(t_0) \rangle \qquad (26)\end{aligned}$$

is the explicit form of the *action principle* for dynamics in the time interval $[t_0, t_1]$.

Note that we are including external forces in the formulation of dynamics and we are not requiring the vanishing of the virtual displacements at the two ends of the time interval.

By using displacements with $\delta\xi(t_0) = 0$ and $\delta\xi(t_1) = 0$ we derive the equality

$$\int_{t_0}^{t_1} \left\langle \frac{\partial L}{\partial q} \circ (\xi, \dot{\xi}) - \frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}) \right), \delta\xi \right\rangle = - \int_{t_0}^{t_1} \langle \varphi, \delta\xi \rangle. \qquad (27)$$

The *Euler-Lagrange equations*

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}) \right) - \frac{\partial L}{\partial q} \circ (\xi, \dot{\xi}) = \varphi \qquad (28)$$

follow. Assuming that these equations are satisfied and using arbitrary displacements we derive the *momentum-velocity relations*

$$\frac{\partial L}{\partial \dot{q}}(\xi(t_0), \dot{\xi}(t_0)) = p_0 \quad \text{and} \quad \frac{\partial L}{\partial \dot{q}}(\xi(t_1), \dot{\xi}(t_0)) = p_1 \qquad (29)$$

from

$$\left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}), \delta\xi \right\rangle (t_1) - \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}), \delta\xi \right\rangle (t_0) = \langle p_1, \delta\xi(t_1) \rangle - \langle p_0, \delta\xi(t_0) \rangle. \qquad (30)$$

We observe that the dynamics $D_{[t_0,t_1]}$ is the set of the solutions of Euler-Lagrange equations to be satisfied inside the time interval $[t_0, t_1]$ plus the momentum velocity relations which have to be satisfied at the two ends of the time interval. The presence of the two momentum velocity relations is the result of using non vanishing virtual displacements at the end points.

4. Phase space trajectories.

The space

$$Ph = Q \times V^* \times V^* \qquad (31)$$

is the *configuration-force-momentum phase space* of the system. Elements (q, f, p) of this space represent the configuration, the external force, and momentum at a given time. A curve

$$(\xi, \varphi, \pi)\colon I \to Ph \qquad (32)$$

defined on an open interval $I \subset \mathbb{R}$ of time is called a *phase space trajectory.* Its component

$$\xi\colon I \to Q \tag{33}$$

is a *configuration space trajectory.* A curve

$$\delta\xi\colon I \to V \tag{34}$$

is a *variation* of the configuration space trajectory ξ. We denote by $\mathcal{P}h$ the space of phase space trajectories (32) and introduce the set $\mathcal{D} \subset \mathcal{P}h$ of trajectories (ξ, φ, π) with the property that for each closed interval $[t_0, t_1]$ included in the domain of definition I the objects

$$(\xi|[t_0,t_1]) = \xi|[t_0,t_1]\,,\ \ (\varphi|[t_0,t_1]) = \varphi|[t_0,t_1]\,,\ \ p_0 = \pi(t_0)\,,\ \ p_1 = \pi(t_1) \tag{35}$$

are in the dynamics $D_{[t_0,t_1]}$.

The action principle for $\mathcal{D}$ states that

$$\begin{aligned}\int_{t_0}^{t_1} \left\langle \frac{\partial L}{\partial q} \circ (\xi,\dot{\xi}) - \frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\partial L}{\partial \dot{q}} \circ (\xi,\dot{\xi})\right), \delta\xi \right\rangle + \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi,\dot{\xi}), \delta\xi \right\rangle (t_1) \\ - \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi,\dot{\xi}), \delta\xi \right\rangle (t_0) = -\int_{t_0}^{t_1} \langle \varphi, \delta\xi \rangle + \langle \pi, \delta\xi\rangle(t_1) - \langle \pi, \delta\xi\rangle(t_0)\end{aligned} \tag{36}$$

for each variation $\delta\xi\colon I \to V$ of the trajectory $\xi\colon I \to Q$ and each time interval $[t_0, t_1]$ contained in the domain of definition I of the phase space trajectory (32).. A trajectory $(\xi, \varphi, \pi)\colon I \to Ph$ is in the set $\mathcal{D}$ if and only if equations

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\partial L}{\partial \dot{q}} \circ (\xi,\dot{\xi})\right) - \frac{\partial L}{\partial q} \circ (\xi,\dot{\xi}) = \varphi \tag{37}$$

and

$$\frac{\partial L}{\partial \dot{q}} \circ (\xi,\dot{\xi}) = \pi \tag{38}$$

are satisfied in the domain of definition I.

The set $\mathcal{D}$ is an alternative representation of dynamics equivalent to the representation in terms of the family D of the solutions of the boundary problems in each time interval $[t_0, t_1] \subset \mathbb{R}$.

5. Intervals as distributions.

We will now show that in a slightly modified formulation of analytical mechanics in a finite time interval $[t_0, t_1]$ this interval is used solely as a domain of integration. Consequently it can be treated as a distribution in $\mathbb{R}$. This fact permits an unified formulation of analytical mechanics in which the passage to the infinitesimal limit is rather simple since an infinitesimal time interval can be introduced as a Dirac distribution.

A *motion* is an equivalence class of pairs $(\xi, [t_0, t_1])$ of a configuration space trajectory $\xi\colon I \to Q$ and an interval $[t_0, t_1] \subset I$. Two pairs $(\xi, [t_0, t_1])$ and $(\xi', [t'_0, t'_1])$ are equivalent if $[t'_0, t'_1] = [t_0, t_1]$ and

$$\int_{t_0}^{t_1} \kappa \circ (\xi', \dot{\xi}', t) = \int_{t_0}^{t_1} \kappa \circ (\xi, \dot{\xi}, t) \tag{39}$$

for each function $\kappa\colon Q \times V \times \mathbb{R} \to \mathbb{R}$. We are presenting an alternative construction of essentially the same set of motions $\Xi(Q|\mathbb{R})$. In this alternative construction of motions restriction of trajectories to the interval $[t_0, t_1]$ is not explicitly used. The interval appears as a domain of integration rather than a set of numbers. We denote by $(\xi|[t_0, t_1])$ the class of $(\xi, [t_0, t_1])$. The construction of functions $K\colon \Xi(Q|\mathbb{R}) \to \mathbb{R}$ which we made in Section 2 is still applicable.

A vector or a *displacement* is an equivalence class of pairs $(\delta\xi, [t_0, t_1])$, where

$$\delta\xi : I \to V \tag{40}$$

is a variation defined on an open interval I containing $[t_0, t_1]$. Pairs $(\delta\xi, [t_0, t_1])$ and $(\delta\xi', [t'_0, t'_1])$ are equivalent if $[t'_0, t'_1] = [t_0, t_1]$ and

$$\frac{\mathrm{d}}{\mathrm{d}s} K(\xi + s\delta\xi'|[t_0, t_1])\big|_{s=0} = \frac{\mathrm{d}}{\mathrm{d}s} K(\xi + s\delta\xi|[t_0, t_1])\big|_{s=0} \tag{41}$$

for each function $K \in \Phi(Q|\mathbb{R})$ and each motion $(\xi|[t_0, t_1])$. We denote by $(\delta\xi|[t_0, t_1])$ the class of $(\delta\xi, [t_0, t_1])$. We have provided an alternative construction of the space $\Delta(Q|\mathbb{R})$.

A covector is an equivalence class of triples $(\varphi, \pi, [t_0, t_1])$, where

$$(\varphi, \pi) : I \to V^* \times V^* \tag{42}$$

is a mapping defined on an open interval I containing $[t_0, t_1]$. Triples $(\varphi, \pi, [t_0, t_1])$ and $(\varphi', \pi', [t'_0, t'_1])$ are equivalent if $[t'_0, t'_1] = [t_0, t_1]$ and

$$\int_{t_0}^{t_1} \Big(\langle \dot{\pi}' - \varphi', \delta\xi \rangle + \langle \pi', \delta\dot{\xi} \rangle \Big) = \int_{t_0}^{t_1} \Big(\langle \dot{\pi} - \varphi, \delta\xi \rangle + \langle \pi, \delta\dot{\xi} \rangle \Big) \tag{43}$$

for each mapping $\delta\xi\colon [t_0, t_1] \to V$. We denote by $(\varphi, \pi|[t_0, t_1])$ the class of $(\varphi, \pi, [t_0, t_1])$. We have described an alternative construction of the space $\Pi(Q|\mathbb{R})$.

The pairing

$$\big\langle\!\big((\varphi, \pi|[t_0, t_1]), (\delta\xi|[t_0, t_1]) \big)\!\big\rangle = \int_{t_0}^{t_1} \Big(\langle \dot{\pi} - \varphi, \delta\xi \rangle + \langle \pi, \delta\dot{\xi} \rangle \Big) \tag{44}$$

between vectors and covectors permits the formulation of the action principle as in the original formulation resulting in the same dynamics in any time interval $[t_0, t_1]$.

6. Unified formulation of analytical mechanics.

We will now describe the unified formulation of analytical mechanics using more general distributions in $\mathbb{R}$ in place of intervals.

A *motion* is an equivalence class of pairs $(\xi, \boldsymbol{c})$ of a configuration space trajectory $\xi\colon I \rightarrow Q$ and a distribution $\boldsymbol{c}$ in $\mathbb{R}$ with compact support $\mathrm{Sup}(\boldsymbol{c}) \subset I$. Two pairs $(\xi, \boldsymbol{c})$ and $(\xi', \boldsymbol{c}')$ are equivalent if $\boldsymbol{c}' = \boldsymbol{c}$ and

$$\int_{\boldsymbol{c}} \kappa \circ (\xi', \dot{\xi}', t) = \int_{\boldsymbol{c}} \kappa \circ (\xi, \dot{\xi}, t) \tag{45}$$

for each function $\kappa\colon Q \times V \times \mathbb{R} \rightarrow \mathbb{R}$. We will denote by $(\xi|\boldsymbol{c})$ the class of $(\xi, \boldsymbol{c})$. The space of motions associated with a distribution $\boldsymbol{c}$ will be denoted by $\Xi(Q|\mathbb{R}, \boldsymbol{c})$ and the space of all motions will be denoted by $\Xi(Q|\mathbb{R})$.

From a differentiable function

$$\kappa\colon Q \times V \times \mathbb{R} \rightarrow \mathbb{R} \tag{46}$$

with arguments $q \in Q$, $\dot{q} \in V$, and $t \in \mathbb{R}$ we construct the function

$$K\colon \Xi(Q|\mathbb{R}) \rightarrow \mathbb{R}\colon (\xi|\boldsymbol{c}) \mapsto \int_{\boldsymbol{c}} \kappa \circ (\xi, \dot{\xi}, t). \tag{47}$$

The symbol $\Phi(Q|\mathbb{R})$ will be used to denote the space of functions on $\Xi(Q|\mathbb{R})$ constructed in this way from functions on $Q \times V \times \mathbb{R}$.

A vector or a *displacement* is an equivalence class of pairs $(\delta\xi, \boldsymbol{c})$, where $\delta\xi$ is a mapping of an open interval $I \subset \mathbb{R}$ in V and $\boldsymbol{c}$ is a distribution with compact support contained in I. Two pairs $(\delta\xi, \boldsymbol{c})$ and $(\delta\xi', \boldsymbol{c}')$ are equivalent if $\boldsymbol{c}' = \boldsymbol{c}$ and

$$\frac{\mathrm{d}}{\mathrm{d}s} K(\xi + s\delta\xi'|\boldsymbol{c})\big|_{s\,=\,0} = \frac{\mathrm{d}}{\mathrm{d}s} K(\xi + s\delta\xi|\boldsymbol{c})\big|_{s\,=\,0} \tag{48}$$

for each function $K \in \Phi(Q|\mathbb{R})$ and each motion $(\xi|\boldsymbol{c})$. The space of displacements associated with a distribution $\boldsymbol{c}$ will be denoted by $\Delta(Q|\mathbb{R}, \boldsymbol{c})$ and the space of all displacements will be denoted by $\Delta(Q|\mathbb{R})$.

A covector is an equivalence class of triples $(\varphi, \pi, \boldsymbol{c})$, where (φ, π) is a mapping of an open interval $I \subset \mathbb{R}$ in $V^* \times V^*$ and $\boldsymbol{c}$ is a distribution with compact support contained in I. Two triples $(\varphi, \pi, \boldsymbol{c})$ and $(\varphi', \pi', \boldsymbol{c}')$ are equivalent if $\boldsymbol{c}' = \boldsymbol{c}$ and

$$\int_{\boldsymbol{c}} \Big(\langle \dot{\pi}' - \varphi', \delta\xi \rangle + \langle \pi', \delta\dot{\xi} \rangle \Big) = \int_{\boldsymbol{c}} \Big(\langle \dot{\pi} - \varphi, \delta\xi \rangle + \langle \pi, \delta\dot{\xi} \rangle \Big) \tag{49}$$

for each mapping $\delta\xi\colon I \rightarrow V$. The interval I is assumed to be the common domain of definition of the mappings (φ, π) and (φ', π'). We will denote by $(\delta\xi|\boldsymbol{c})$ the class of $(\delta\xi, \boldsymbol{c})$ and by $(\varphi, \pi|\boldsymbol{c})$ the class of (φ, π). The space of covectors associated with a distribution $\boldsymbol{c}$ will be denoted by $\Pi(Q|\mathbb{R}, \boldsymbol{c})$ and the space of all covectors will be denoted by $\Pi(Q|\mathbb{R})$.

The originally introduced spaces $\Xi(Q|\mathbb{R})$, $\Delta(Q|\mathbb{R})$, and $\Pi(Q|\mathbb{R})$ are subsets of the new constructions. The set $\Xi(Q|\mathbb{R},\boldsymbol{c})$ is an affine space and the vector space $\Delta(Q|\mathbb{R},\boldsymbol{c})$ is its natural model space.

The space $\Pi(Q|\mathbb{R},\boldsymbol{c})$ is a dual of the space $\Delta(Q|\mathbb{R},\boldsymbol{c})$. The pairing

$$\boldsymbol{\langle}(\varphi,\pi|\boldsymbol{c}),(\delta\xi|\boldsymbol{c})\boldsymbol{\rangle}_{\boldsymbol{c}} = \int_{\boldsymbol{c}} \Big(\langle\dot{\pi}-\varphi,\delta\xi\rangle + \langle\pi,\delta\dot{\xi}\rangle\Big) \tag{50}$$

between vectors and covectors permits the formulation of the virtual action principle.

Let

$$L\colon Q\times V\to\mathbb{R} \tag{51}$$

be the *Lagrangian* of the system. The *action* is the function

$$W\colon \Xi(Q|\mathbb{R})\to\mathbb{R}\colon (\xi|\boldsymbol{c})\mapsto \int_{\boldsymbol{c}} L\circ(\xi,\dot{\xi}). \tag{52}$$

The principle of virtual action is included in the definition of the *dynamics*. The *dynamics associated with the distribution* $\boldsymbol{c}$ is the set

$$\begin{aligned} D_{\boldsymbol{c}} = \Big\{&((\xi|\boldsymbol{c}),(\varphi,\pi|\boldsymbol{c}))\in\Xi(Q|\mathbb{R},\boldsymbol{c})\times\Pi(Q|\mathbb{R},\boldsymbol{c});\\ &\forall_{(\delta\xi|\boldsymbol{c})\in\Delta(Q;\mathbb{R}|\boldsymbol{c})}\ \boldsymbol{\langle}(\varphi,\pi|\boldsymbol{c}),(\delta\xi|\boldsymbol{c})\boldsymbol{\rangle}_{\boldsymbol{c}} = \mathrm{D}W((\xi|\boldsymbol{c}),(\delta\xi|\boldsymbol{c}))\Big\}. \end{aligned} \tag{53}$$

The *dynamics* is the family

$$D = \bigcup_{\boldsymbol{c}\in\mathsf{C}\mathbb{R}} D_{\boldsymbol{c}} \tag{54}$$

where $\mathsf{C}\mathbb{R}$ is the set of the distributions in $\mathbb{R}$ with compact support.

The distribution $\boldsymbol{c}$ is usually derived from an interval $[t_0,t_1]\subset\mathbb{R}$ or is the Dirac delta at $t\in\mathbb{R}$. If $\boldsymbol{c}=[t_0,t_1]$, then the original formulation of analytical mechanics in a finite time interval is obtained. If $\boldsymbol{c}=\delta(t)$, then the infinitesimal limit of the dynamics, which in our terminology is called Lagrangian formulation, is obtained.

7. Infinitesimal limit of the dynamics in terms of the Dirac delta.

Let the distribution $\boldsymbol{c}$ be the Dirac distribution $\boldsymbol{c}=\delta(t)$. The construction of motions, displacements and covectors is based on the expressions

$$\int_{\delta(t)} (\kappa\circ(\xi,\dot{\xi},t)) = (\kappa\circ(\xi,\dot{\xi},t))(t), \tag{55}$$

$$\frac{\mathrm{d}}{\mathrm{d}s}K(\xi+s\delta\xi|\delta(t))\Big|_{s\,=\,0} = (\mathrm{D}\kappa\circ(\xi,\dot{\xi},t,\delta\xi,\delta\dot{\xi},0))(t), \tag{56}$$

and

$$\int_{\delta(t)} \Big(\langle\dot{\pi}-\varphi,\delta\xi\rangle + \langle\pi,\delta\dot{\xi}\rangle\Big) = \Big(\langle\dot{\pi}-\varphi,\delta\xi\rangle + \langle\pi,\delta\dot{\xi}\rangle\Big)(t). \tag{57}$$

It follows that a motion $(\xi|\delta(t))$ is represented by the pair

$$(q, \dot{q}) = (\xi(t), \dot{\xi}(t)) \in Q \times V, \tag{58}$$

a displacement is represented by the pair

$$(\delta q, \delta\dot{q}) = (\delta\xi(t), \delta\dot{\xi}(t)) \in V \times V, \tag{59}$$

and a covector is represented by the pair

$$(r, p) = (\dot{\pi}(t) - \varphi(t), \pi(t)) \in V^* \times V^*, \tag{60}$$

with the pairing

$$\langle\!\langle (r, p), (\delta q, \delta\dot{q}) \rangle\!\rangle_{\delta(t)} = \langle r, \delta q\rangle + \langle p, \delta\dot{q}\rangle. \tag{61}$$

The dynamics associated with the Dirac distribution $\delta(t)$, obtained from the definition (53), is the infinitesimal dynamics

$$\begin{aligned} D_{\delta(t)} = \Big\{ &(q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*; \\ &\qquad\qquad \forall_{(\delta q, \delta\dot{q}) \in V \times V}\ \langle r, \delta q\rangle + \langle p, \delta\dot{q}\rangle = \mathrm{D}L(q, \dot{q}, \delta q, \delta\dot{q}) \Big\} \\ = \Big\{ &(q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*;\ \frac{\partial L(q,\dot{q})}{\partial q} = r,\ \frac{\partial L(q,\dot{q})}{\partial \dot{q}} = p \Big\}. \end{aligned} \tag{62}$$

Note that the constructed spaces $\Xi(Q|\mathbb{R}, \delta(t))$, $\Delta(Q|\mathbb{R}, \delta(t))$, and $\Pi(Q|\mathbb{R}, \delta(t))$ and also the infinitesimal dynamics $D_{\delta(t)}$ of an autonomous system with a time independent Lagrangian are the same for each $t \in \mathbb{R}$. We will replace the symbol $\delta(t)$ with δ.

We show that a trajectory $(\xi, \varphi, \pi)\colon I \to Ph$ is in the set $\mathcal{D}$ if and only if

$$(\xi, \pi, \dot{\xi}, \dot{\pi} - \varphi)(t) \in D_\delta \tag{63}$$

for each $t \in I$. The proof is based on the equality

$$\int_{t_0}^{t_1} \left(\left\langle \frac{\partial L}{\partial q} \circ (\xi, \dot{\xi}), \delta\xi \right\rangle + \left\langle \frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}), \delta\dot{\xi} \right\rangle \right) = \int_{t_0}^{t_1} \Big(\langle \dot{\pi} - \varphi, \delta\xi\rangle + \langle \pi, \delta\dot{\xi}\rangle \Big) \tag{64}$$

which is a version of the principle of virtual action. If a curve $(\xi, \varphi, \pi)\colon I \to Ph$ is in $\mathcal{D}$ and $\delta\xi\colon I \to V$ is an arbitrary variation, then the equality (64) holds for all intervals $[t_0, t_1] \subset I$. It follows that the equality

$$\begin{aligned} \left\langle \frac{\partial L}{\partial q}(\xi(t), \dot{\xi}(t)), \delta\xi(t) \right\rangle + \left\langle \frac{\partial L}{\partial \dot{q}}(\xi(t), \dot{\xi}(t)), \delta\dot{\xi}(t) \right\rangle &= \langle \dot{\pi}(t) - \varphi(t), \delta\xi(t)\rangle \\ &\quad + \langle \pi(t), \delta\dot{\xi}(t)\rangle \end{aligned} \tag{65}$$

holds at each $t \in I$. This implies that $(\xi, \pi, \dot{\xi}, \dot{\pi} - \varphi)(t) \in D_\delta$ for each $t \in I$ thanks to the arbitrariness of the vectors $\delta\xi(t)$ and $\delta\dot{\xi}(t)$. Conversely if $(\xi, \varphi, \pi) \colon I \to Ph$ is such that $(\xi, \pi, \dot{\xi}, \dot{\pi} - \varphi)(t) \in D_\delta$ for each $t \in I$, then the equality (65) holds in I with an arbitrary variation $\delta\xi \colon I \to V$. It follows by integration that (ξ, φ, π) satisfies the principle of virtual action for each time interval $[t_0, t_1] \subset I$. Hence, the trajectory (ξ, φ, π) is in the set $\mathcal{D}$.

Equations

$$\frac{\partial L}{\partial q} \circ (\xi, \dot{\xi}) = \dot{\pi} - \varphi \tag{66}$$

and

$$\frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi}) = \pi \tag{67}$$

are obtained for the phase space trajectories. These equations will be called the *Lagrange equations.* The first order Lagrange equations (66) and (67) are equivalent to the second order Euler-Lagrange equations (37) together with (38) or (67). The equivalence is established by using the equation

$$\frac{\mathrm{d}}{\mathrm{d}t}\left(\frac{\partial L}{\partial \dot{q}} \circ (\xi, \dot{\xi})\right) = \dot{\pi}. \tag{68}$$

8. Hamiltonian formulation of dynamics.

From a Lagrangian $L \colon Q \times V \to \mathbb{R}$ we derive the *energy function*

$$E \colon Q \times V^* \times V \to \mathbb{R} \colon (q, p, \dot{q}) \mapsto \langle p, \dot{q} \rangle - L(q, \dot{q}) \tag{69}$$

and associate with this function the family (E, η) of functions

$$E(q, p, \,\cdot\,) \colon V \to \mathbb{R} \tag{70}$$

on fibres of the projection

$$\eta \colon Q \times V^* \times V \to Q \times V^*. \tag{71}$$

We show that the energy function can be used to generate the infinitesimal dynamics D_δ. The family (E, η) generates the set

$$D = \Big\{(q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*;\ \exists_{v \in V}\ \forall_{(\delta q, \delta p, \delta v) \in V \times V^* \times V} \\ \langle r, \delta q \rangle - \langle \delta p, \dot{q} \rangle = -\mathrm{D}E(q, p, v, \delta q, \delta p, \delta v)\Big\}. \tag{72}$$

The use of families of functions as generating objects for sets is described in [3].

The set (72) is obtained by projecting the set

$$\overline{D} = \Big\{(q, p, \dot{q}, r, v) \in Q \times V^* \times V \times V^* \times V;\ \forall_{(\delta q, \delta p, \delta v) \in V \times V^* \times V} \\ \langle r, \delta q \rangle - \langle \delta p, \dot{q} \rangle = -\mathrm{D}E(q, p, v, \delta q, \delta p, \delta v)\Big\} \tag{73}$$

onto $Q \times V^* \times V \times V^*$. By setting $\delta q = 0$ and $\delta v = 0$ and using

$$\mathrm{D}E(q,p,v,\delta q,\delta p,\delta v) = \langle \delta p, v\rangle + \langle p, \delta v\rangle - \mathrm{D}L(q,v,\delta q,\delta v) \tag{74}$$

we derive from the definition of $\overline{D}$ the equality $v = \dot q$. The set $\overline{D}$ reduces to

$$\begin{aligned}\overline{D} = \Big\{&(q,p,\dot q,r,v) \in Q \times V^* \times V \times V^* \times V;\ v = \dot q \\ &\forall_{(\delta q,\delta \dot q)\in V\times V}\ \langle r,\delta q\rangle + \langle p,\delta \dot q\rangle = \mathrm{D}L(q,\dot q,\delta q,\delta \dot q)\Big\}.\end{aligned} \tag{75}$$

It projects onto the infinitesimal dynamics

$$\begin{aligned}D_\delta = \Big\{&(q,p,\dot q,r) \in Q \times V^* \times V \times V^*; \\ &\forall_{(\delta q,\delta \dot q)\in V\times V}\ \langle r,\delta q\rangle + \langle p,\delta \dot q\rangle = \mathrm{D}L(q,\dot q,\delta q,\delta \dot q)\Big\}.\end{aligned} \tag{76}$$

Hence $D = D_\delta$. The appearance of the *Hamiltonian pairing*

$$\langle\!\langle (q,p,\dot q,r),(q,p,\delta q,\delta p)\rangle\!\rangle^{\mathsf{H}} = \langle r,\delta q\rangle - \langle \delta p,\dot q\rangle \tag{77}$$

in the formula (72) means that the family (E,η) represents a *Hamiltonian generating object.* The *Lagrangian pairing*

$$\langle\!\langle (q,p,\dot q,r),(q,\dot q,\delta q,\delta \dot q)\rangle\!\rangle^{\mathsf{L}} = \langle r,\delta q\rangle + \langle p,\delta \dot q\rangle \tag{78}$$

was used to generate the dynamics D_δ from the Lagrangian.

The set

$$\begin{aligned}Cr(E,\eta) &= \Big\{(q,p,\dot q) \in Q \times V^* \times V;\ \forall_{\delta \dot q \in V}\ \mathrm{D}E(q,p,\dot q,0,0,\delta \dot q) = 0\Big\} \\ &= \Big\{(q,p,\dot q) \in Q \times V^* \times V;\ p = \frac{\partial L}{\partial \dot q}(q,\dot q)\Big\}\end{aligned} \tag{79}$$

is the *critical set* of the family (E,η). The critical set is the graph of the *Legendre mapping*

$$\lambda\colon Q \times V \to V^*\colon (q,\dot q) \mapsto \frac{\partial L}{\partial \dot q}(q,\dot q). \tag{80}$$

A Lagrangian is said to be *hyperregular* if the critical set is the image of a section

$$\sigma\colon Q \times V^* \to Q \times V^* \times V \tag{81}$$

of the projection η. This is the case if for each q the mapping

$$\lambda(q,\,\cdot\,)\colon V \to V^* \tag{82}$$

is invertible. It is convenient to use the mapping

$$\rho\colon Q \times V^{*} \to V\colon (q,p) \mapsto \lambda(q,\,\cdot\,)^{-1}(p). \tag{83}$$

The section σ is then defined by

$$\sigma(q,p) = (q,p,\rho(q,p))\,. \tag{84}$$

If the Lagrangian is hyperregular, then the *Hamiltonian*

$$H = E \circ \sigma\colon Q \times V^{*} \to \mathbb{R}\colon (q,p) \mapsto E(q,p,\rho(q,p)) \tag{85}$$

is introduced. The passage from the Lagrangian L to the Hamiltonian H is the *Legendre transformation of analytical mechanics*.

Assuming that the Lagrangian is hyperregular we show that the dynamics is generated by the Hamiltonian. By setting $\delta q = 0$ and $\delta p = 0$ in (73) we derive the equality

$$\forall_{\delta v \in V}\ \mathrm{D}E(q,p,v,0,0,\delta v) = 0. \tag{86}$$

It follows that the set $\overline{D}$ is included in the set

$$\left\{(q,p,\dot{q},r,v) \in Q \times V^{*} \times V \times V^{*} \times V;\ (q,p,v) \in Cr(E,\eta)\right\}. \tag{87}$$

Hence,

$$\begin{aligned}\overline{D} = \Big\{(q,p,\dot{q},r,v) \in Q \times V^{*} \times V \times V^{*} \times V;\ &\forall_{(\delta q,\delta p,\delta v) \in V \times V^{*} \times V} \\ &\langle r,\delta q\rangle - \langle \delta p,\dot{q}\rangle = -\mathrm{D}E(q,p,\rho(q,p),\delta q,\delta p,\delta v)\Big\}.\end{aligned} \tag{88}$$

If $(q,p,\dot{q},r,v) \in \overline{D}$, then the equality

$$\langle r,\delta q\rangle - \langle \delta p,\dot{q}\rangle = -\mathrm{D}E(q,p,\rho(q,p),\delta q,\delta p,\delta v) \tag{89}$$

holds for all displacements $(\delta q,\delta p,\delta v)$ and in particular for

$$(\delta q,\delta p,\delta v) = (\delta q,\delta p,\mathrm{D}\rho(q,p,\delta q,\delta p)). \tag{90}$$

The equality

$$\mathrm{D}H(q,p,\delta q,\delta p) = \mathrm{D}E(q,p,\rho(q,p),\delta q,\delta p,\mathrm{D}\rho(q,p,\delta q,\delta p)) \tag{91}$$

implies that $(q,p,\dot{q},r,v)$ is in the set

$$\begin{aligned}\Big\{&(q,p,\dot{q},r,v) \in Q \times V^{*} \times V \times V^{*} \times V; \\ &\qquad \forall_{(\delta q,\delta p) \in V \times V^{*}}\ \langle r,\delta q\rangle - \langle \delta p,\dot{q}\rangle = -\mathrm{D}H(q,p,\delta q,\delta p)\Big\}.\end{aligned} \tag{92}$$

The inclusion relation

$$D_\delta \subset \widetilde{D} = \Big\{ (q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*;\ \forall_{(\delta q, \delta p) \in V \times V^*} \\ \langle r, \delta q \rangle - \langle \delta p, \dot{q} \rangle = -\mathrm{D}H(q, p, \delta q, \delta p) \Big\}. \tag{93}$$

follows. An arbitrary displacement $(\delta q, \delta p, \delta v)$ can be presented as the sum

$$(\delta q, \delta p, \delta v) = (\delta q, \delta p, \mathrm{D}\rho(q, p, \delta q, \delta p)) + (0, 0, \delta v - \mathrm{D}\rho(q, p, \delta q, \delta p)). \tag{94}$$

If $(q, p, \dot{q}, r) \in \widetilde{D}$, then the equality

$$\langle r, \delta q \rangle - \langle \delta p, \dot{q} \rangle = -\mathrm{D}E(q, p, \rho(q, p), \delta q, \delta p, \mathrm{D}\rho(q, p, \delta q, \delta p)) \tag{95}$$

holds for all displacements $(\delta q, \delta p)$. The equality

$$\mathrm{D}E(q, p, \rho(q, p), 0, 0, \delta v - \mathrm{D}\rho(q, p, \delta q, \delta p)) = 0 \tag{96}$$

holds as well since $(q, p, \rho(q, p))$ is in the critical set. It follows that $(q, p, \dot{q}, r, \rho(q, p))$ is in $\overline{D}$ and $(q, p, \dot{q}, r)$ is in D_δ. We have found that

$$D_\delta = \Big\{ (q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*; \\ \forall_{(\delta q, \delta p) \in V \times V^*}\ \langle r, \delta q \rangle - \langle \delta p, \dot{q} \rangle = -\mathrm{D}H(q, p, \delta q, \delta p) \Big\}. \tag{97}$$

This expression for the infinitesimal dynamics is the *Hamiltonian representation* of dynamics. Equations

$$\frac{\partial H}{\partial q} \circ (\xi, \pi) = \varphi - \dot{\pi}\ , \quad \frac{\partial H}{\partial p} \circ (\xi, \pi) = \dot{\xi} \tag{98}$$

derived for phase space trajectories (ξ, φ, π) from the above Hamiltonian representation of D_δ are the *canonical Hamilton equations.*

We use the following example to illustrate the theory which we have formulated.

EXAMPLE 2. We will describe the dynamics of the system whose statics was studied in Example 1. The Lagrangian of the system is the function

$$L\colon Q \times V \to \mathbb{R}\colon (q, \dot{q}) \mapsto \frac{m}{2}\langle g(\dot{q}), \dot{q} \rangle - \frac{k}{2}\langle g(q - q_0), q - q_0 \rangle, \tag{99}$$

composed of the *kinetic energy*

$$T\colon V \to \mathbb{R}\colon \dot{q} \mapsto \frac{m}{2}\langle g(\dot{q}), \dot{q} \rangle \tag{100}$$

and the *potential energy*

$$U\colon Q \to \mathbb{R}\colon q \mapsto \frac{k}{2}\langle g(q - q_0), q - q_0 \rangle. \tag{101}$$

The *dynamics in the interval* $[t_0, t_1]$ is the set

$$\begin{aligned} D_{[t_0,t_1]} = \Big\{ &((\xi|[t_0,t_1]),(\varphi|[t_0,t_1],p_0,p_1)) \in \Xi(Q|\mathbb{R},[t_0,t_1]) \times \Pi(Q;\mathbb{R}|[t_0,t_1]); \\ &mg\circ\ddot{\xi} + k(\xi - q_0) = \varphi,\ mg(\dot{\xi}(t_0)) = p_0,\ mg(\dot{\xi}(t_1)) = p_1 \Big\}. \end{aligned} \tag{102}$$

The Euler-Lagrange equations to be satisfied in the interval $[t_0, t_1]$ are

$$mg\circ\ddot{\xi} + k(\xi - q_0) = \varphi, \tag{103}$$

with

$$\ddot{\xi}\colon [t_0,t_1] \to V\colon t \mapsto \frac{\mathrm{d}^2\xi(t)}{\mathrm{d}t^2}, \tag{104}$$

and the relations

$$mg(\dot{\xi}(t_0)) = p_0, \quad \text{and} \quad mg(\dot{\xi}(t_1)) = p_1, \tag{105}$$

are the momentum-velocity relations.

The familiar equations

$$mg\circ\ddot{\xi} + k(\xi - q_0) = \varphi \tag{106}$$

and

$$mg\circ\dot{\xi} = \pi \tag{107}$$

are derived for phase space trajectories (ξ, φ, π) from the Lagrangian (99).

The infinitesimal dynamics in the Lagrangian formulation has the following expression.

$$D_\delta = \big\{(q,p,\dot{q},r) \in Q \times V^* \times V \times V^*;\ k(q - q_0) = -r,\ mg(\dot{q}) = p\big\}. \tag{108}$$

We recall that we set $r = \dot{p} - f$.

The Lagrange equations have the form

$$k(\xi - q_0) = \varphi - \dot{\pi}, \tag{109}$$

and

$$mg\circ\dot{\xi} = \pi. \tag{110}$$

The mapping

$$E\colon Q \times V^* \times V \to \mathbb{R}\colon (q,p,\dot{q}) \mapsto \langle p,\dot{q}\rangle - \frac{m}{2}\langle g(\dot{q}),\dot{q}\rangle + \frac{k}{2}\langle g(q-q_0), q-q_0\rangle \tag{111}$$

is the energy function. The set

$$Cr(E,\eta) = \big\{(q,p,\dot{q}) \in Q \times V^* \times V;\ p = mg(\dot{q})\big\} \tag{112}$$

is the critical set of the family (E, η), where η is the projection introduced in the preceding section. It is the graph of the Legendre mapping

$$\lambda \colon Q \times V \to V^* \colon (q, \dot{q}) \mapsto mg(\dot{q}). \tag{113}$$

The mapping

$$\lambda(q, \,\cdot\,) \colon V \to V^* \colon \dot{q} \mapsto mg(\dot{q}) \tag{114}$$

is invertible. Its inverse is the mapping

$$\lambda(q, \,\cdot\,)^{-1} \colon V^* \to V \colon p \mapsto \frac{1}{m} g^{-1}(p). \tag{115}$$

Hence, the Lagrangian (99) is hyperregular and the Hamiltonian function

$$H \colon Q \times V^* \to \mathbb{R} \colon (q, p) \mapsto \frac{1}{2m} \langle p, g^{-1}(p) \rangle + \frac{k}{2} \langle g(q - q_0), q - q_0 \rangle \tag{116}$$

is derived from the family (E, η).

In the Hamiltonian formulation the infinitesimal dynamics has the following expression.

$$D_\delta = \left\{ (q, p, \dot{q}, r) \in Q \times V^* \times V \times V^*; \; k(q - q_0) = -r, \; \dot{q} = \frac{1}{m} g^{-1}(p) \right\}. \tag{117}$$

The Hamilton equations for phase space trajectories (ξ, φ, π) have the form

$$k(\xi - q_0) = \varphi - \dot{\pi}, \tag{118}$$

and

$$\dot{\xi} = \frac{1}{m} g^{-1} \circ \pi. \tag{119}$$

▲

REFERENCES

[1] W. M. Tulczyjew, *The origin of variational principles*, in the volume Classical and quantum integrability (Warsaw, 2001), 41–75, Banach Center Publ., **59**, Polish Acad. Sci., Warsaw, 2003.

[2] G. Marmo, W. M. Tulczyjew, P. Urbański, *Dynamics of autonomous systems with external forces*, Acta Physica Polonica B, **33** (2002), 1181–1240.

[3] W. M. Tulczyjew, P. Urbański, *Homogeneous Lagrangian systems*, in *Gravitation, Electromagnetism and Geometric Structures*, Pitagora Editrice, 1996. See also arXiv: math-ph/0404075.